



SYNTHESIS AND CHARACTERIZATION OF 2-(1H-BENZIMIDAZOL-2-YL-SULFANYL) ACETO HYDRAZIDE

RAMESH DHANI

Department of Pharmaceutical Chemistry, Oil Technological Research Institute, Jawaharlal
Nehru Technological University Anantapur, Andhra Pradesh, India

Abstract

Accepted Date:

08/10/2012

Publish Date:

27/10/2012

Keywords

Benzimidazole

Imidazole

N-Hexane

Ethyl acetate

Chloroform

Methanol

An important feature of modern pharmaceutical chemistry is the introduction of more refined and sensitive methods of physicochemical analysis such as Spectroscopy and chromatography that enable one to assay the quality and quantity of the drugs more accurately with the smallest consumption of the analyze, reagent and time. Diversity of biological response profile has attracted considerable interest of several researchers across the globe to explore this skeleton for its assorted therapeutic significance. By using different synthetic methods new benzimidazole derivatives were synthesized and further Melting points were determined by using Precision melting point apparatus in open capillaries and are uncorrected. The purity of the compounds was checked by TLC on silica gel G plates using n-Hexane, ethyl acetate (1:3) and methanol: chloroform (1:9) solvent system. The synthesized benzimidazole derivatives were characterized by IR spectral analysis. Benzimidazole is a lead nucleus for future developments to get effective compounds.

Corresponding Author

Mr. Ramesh Dhani

INTRODUCTION

Pharmaceutical chemistry is a science that makes use of general laws of chemistry to study drugs, i.e. their preparation, chemical nature, composition, structure, influence on an organism and studies of the physical and chemical properties of drugs, the methods of quality control and the conditions of their storage. Pharmaceutical chemistry occupies the most important place among the related sciences e.g. Drug technology, Toxicological chemistry, Pharmacognosy, and the organization of the pharmacy.


Heterocyclic chemistry is the branch of chemistry dealing with the synthesis, properties, and applications of heterocyclics. Heterocyclic chemistry comprises at least half of all organic chemistry research worldwide. In particular, heterocyclic structures form the basis of many pharmaceutical, agrochemical and veterinary products.

The benzimidazole contains a phenyl ring fused to an imidazole ring, as indicated in the structure of benzimidazole. The important group of substances has found practical application in a number of fields.

Recently in benzimidazole chemistry has been revived somewhat by the discovery that the 5, 6-dimethyl benzimidazole moiety is a part of the chemical structure of vitamin B12.

MATERIALS & METHODS

1. Synthesis of 2-mercapto benzimidazole




A mixture of 10.8gm (0.1mole) of o-phenylenediamine, 5.65 gm (0.1mole) of potassium hydroxide and 7.67 gm (0.1mole, 6.19ml) of carbon disulfide, 100ml of 95% ethanol and 15 ml of water was taken in a 500ml round bottom flask heated under reflux for three hours. Then 1-1.5 gm of charcoal was added cautiously and the mixture is further heated at the reflux for 10 minutes, the charcoal is removed by filtration. The filtrate is heated to 60-70°C, 100ml of warm water is added, and acidified with dilute acetic acid with good stirring. The product separated as glistening white crystals, and the mixture is placed in a refrigerator for three hours to complete the

crystallization. The product is collected on a Buckner funnel and dried over night at 40°C. The dried product is recrystallised by ethanol the yield is 8.5gm (73%) melting point is 300-305°C.


2. Synthesis of ethyl (1H-benzimidazol-2-yl-sulfanyl) acetate- I

A stirred mixture containing 4.5gm of (0.03mole) of 2-mercaptobenzimidazole, 60ml of ethanol and 1.68gm of (0.03mole) potassium hydroxide was added and heated at 78-80°C for 10-minutes. Then ethyl chloro acetate (3.66ml, 0.03mol) was added in one portion, an exothermic reaction set in causing a temperature rise from 30-40°C. After stirring at 25-30°C for 18-hours, the reaction mixture was added to 100gm of ice-water and stirred for 30-minutes at 0-10°C. The precipitate was collected by filtration washed with water until free of chloride and air dried at 50°C and recrystallised by water the yield is 6 gm (62.25%). melting point is 105°C.



3. Synthesis of 2-(1H-benzimidazol-2-ylsulfanyl) acetohydrazide -II

The mixture of 2-carboxy ethyl thio 1H-benzimidazole 4gm (0.004mole) and Hydrazine hydrate 6ml (0.01mole) are mixed well in a RBF and heated on water bath for 10 min. then dissolved in 60 ml ethanol, the reaction mixture is heated with reflux the reaction mixture is heated with reflux condenser for six hours, cooled to room temperature and the reaction mixture was added to 100gm of ice-water, and kept aside for the crystallization. The colorless crystals are collected by filtration, and recrystallised from water. Melting point is 180-185°C; the yield is 60-70%.



RESULTS AND DISCUSSION

The present study explains the synthesis and characterization of benzimidazol derivative i.e., 2-(1H-benzimidazol-2-yl-sulfanyl) acetohydrazide. The synthesized benzimidazol derivative characterized by IR spectral studies. By this studies find the structural activity relationship (SAR) and to optimize the structure. The purity of the synthesized benzimidazol derivative was checked by thin layer chromatography (TLC) and R_f value was recorded.

IR Spectral Data:

The IR spectrum of the compound IV was recorded on FTIR spectrometer by KBr method. The FTIR spectra from the figure.1 and the table.4 shows bands at 3352 cm^{-1} , 3050 cm^{-1} , 1666 cm^{-1} , 1614 cm^{-1} and 721 cm^{-1} , 672 cm^{-1} corresponds to -NHNH_2 , (aromatic ring), -C=O of amide, -C=N- , C-S-C , (C-S) respectively.

CONCLUSION

By this study concluded that to find the structure-activity relationship (SAR) and to

optimize the structure of the synthesized new benzimidazol derivative i.e., 2-(1H-benzimidazol-2-yl-sulfanyl) acetohydrazide. The compound was characterized by IR spectral studies, the purity of the compound was checked by TLC and it produces good yield. The compound was confirmed by physicochemical and spectral analysis.

ACKNOWLEDGEMENT

I am indebted to my parents and my sister for their inspiration and encouragement given to me during this work with deep appreciation for their determination and enthusiasm at each and every front of my life to transform my dreams into reality.

I express my whole hearted indebtedness and heartfelt thanks to my school teacher's and to my professor's of JNTUA-OTRI Anantapur. I am very thankful and prevail age to my deep sense of gratitude to **Dr.**

Bishwaranjan Behra, M. Pharm., Ph.D.

PHYSICO-CHEMICAL ANALYSIS

Table 1

S.no	2-mercapto benzimidazole	
1.	Mol. Formula	C ₇ H ₆ N ₂ S
2.	Melting Point	300-305°C
3.	% Yield	73%
4.	Solvent system used	hexane: ethyl acetate (1:3)

Table 2

S.no	Ethyl (1H-benzimidazol-2-yl-sulfanyl) acetate	
1.	Mol. Formula	C ₁₁ H ₁₂ O ₂ N ₂ S
2.	Melting Point	105°C
3.	% Yield	62.25 %
4.	Solvent system used	hexane: ethyl acetate (1:3)

Table 3

Sr. no	2-(1H-benzimidazol-2-yl-sulfanyl)acetohydrazide	
1.	Mol. Formula	C ₈ H ₁₁ ON ₄ S
2.	Melting Point	180-185°C
3.	% Yield	60-70%
4.	Solvent system used	hexane: ethyl acetate (1:3)

Table 4

IR values of 2-(1H-benzimidazol-2-yl-sulfanyl) acetohydrazide

Functional group assigned	Group frequencies (cm ⁻¹)
-NHNH2	3352
(aromatic ring),	3050
-C=O of amide	1666
(-C=N-)	1614
C-S-C	721
(C-S)	672




Figure 1 IR Spectra of 2-(1H-benzimidazol-2-yl-sulfanyl) acetohydrazide

REFERENCES

1. Burger A: Comprehensive Medicinal Chemistry. 1990; 23: 22.
2. Camille G and Wermuth: The Practice of Medicinal Chemistry. Elsevier. 2001; 2: 29.
3. Martin Y.C: Modern Drug Research, Paths to Better and Safer Drugs, ed 5, New York, 1989: 243-73.
4. Lindberg P and Mitscher LA: European Journal of Medicinal Chemistry.1998; 6: 243-73.
5. John B Wright: The chemistry of benzimidazoles. Chemical Reviews.1951; 3 (3): 397-541.
6. Day AR: Electronic mechanism of organic synthesis. 1951; 2: 242-243.
7. Preston PN: Synthesis Reactions and Spectroscopic properties of benzimidazoles. Chemical Reviews. 1974; 15: 279-314.
8. Wagner EC and Millett WH: Synthesis of Benzimidazole. Chemistry of Org Synthesis. 1943; 2: 65.
9. Shriner RL and Robert W Upson: Synthesis of bis-benzimidazoles from diabasic acids. Journal of Organic Chemistry. 1941; 63(8): 2277.
10. Walther R Von and Kessler A: Journal of Prakt Chemistry. 1906; 174 : 245.

